

Joint work with Kyle Genova, Frans Schalekamp, and Anke van Zuylen

Outline

- A brief intro to the TSP
- A standard TSP linear program: the Subtour LP
 - Experimental analysis
 - Theoretical analysis: an outstanding open question
- A related question: the Boyd-Carr conjecture and its proof
- Some conjectures and more experiments

The Traveling Salesman Problem

The most famous problem in discrete optimization: Given ncities and the cost c(i,j) of traveling from city i to city j, find a minimum-cost tour that visits each city exactly once.

We assume costs are symmetric (c(i,j)=c(j,i) for all i,j) and obey the triangle inequality $(c(i,j) \le c(i,k) + c(k,j)$ for all i,j,k).

120 city tour of West Germany due to M. Grötschel (1977)



A 15112 city instance solved by Applegate, Bixby, Chvátal, and Cook (2001)



A 24978 city instance from Sweden solved by Applegate, Bixby, Chvátal, Cook, and Helsgaun (2004)





The Dantzig-Fulkerson-Johnson Method

- G=(V,E) is a complete graph on |V| = n vertices
- c(e)=c(i,j) is the cost of traveling on edge
 e=(i,j)
- x(e) is a decision variable indicating if edge e is used in the tour, $0 \le x(e) \le 1$
- Solve linear program; if x(e) forms integer tour, stop, else find a *cutting plane*

The linear program

Minimize
$$\sum_{e \in E} c(e)x(e)$$

subject to

$$\sum_{e \in \delta(v)} x(e) = 2 \qquad \forall v \in V$$

 $0 \le x(e) \le 1 \qquad \forall e \in E$



Fractional 2-matchings



Fractional (basic) solutions have components that are cycles of size at least 3 with x(e)=1 or odd cycles with x(e)=1/2 connected by paths with x(e)=1



Integer solutions have components with cycles of size at least 3; sometimes called *subtours*

"Loop conditions"

Dantzig, Fulkerson, and Johnson added constraints to eliminate subtours as they occurred; these now called "subtour elimination constraints".



Subtour LP

$$\begin{array}{ll} \text{Minimize } \sum_{e \in E} c(e) x(e) \\ \text{subject to} \\ & \sum_{e \in \delta(v)} x(e) = 2 \qquad \forall v \in V \\ & \sum_{e \in \delta(S)} x(e) \geq 2 \qquad \forall S \subseteq V, |S| \geq 2 \\ & 0 \leq x(e) \leq 1 \qquad \forall e \in E \end{array}$$

Equivalent constraints

Equivalently can write subtour elimination constraints to express no cycles in any strict subset:

$$\sum_{e \in E(S)} x(e) \le |S| - 1 \qquad \forall S \subset V, |S| \ge 2$$



Subtour LP



How strong is the Subtour LP bound?

Johnson, McGeoch, and Rothberg (1996) and Johnson and McGeoch (2002) report experimentally that the Subtour LP is very close to the optimal.

Random Uniform Euclidean			TSPLIB				
Name	%Gap	Opttime	HKtime	Name	%Gap	Opttime	HKtime
E1k.0	0.77	1406	2.13	dsj1000	0.61	410	3.68
E1k.1	0.64	3855	2.15	pr1002	0.89	34	2.40
E1k.2	0.72	1211	2.02	si1032	0.08	25	11.32
E1k.3	0.62	956	1.92	u1060	0.65	571	3.62
E1k.4	0.69	330	1.69	vm1084	1.33	605	2.40
E1k.5	0.59	233	2.42	pcb1173	0.96	468	1.70
E1k.6	0.79	2940	1.67	d1291	1.18	27394	4.54
E1k.7	0.94	8003	1.95	rl1304	1.55	189	4.08
E1k.8	1.01	4347	1.65	rl1323	1.65	3742	4.49
E1k.9	0.61	189	2.14	nrw1379	0.43	578	2.40
E3k.0	0.71	533368	9.57	fl1400	1.74	1549	9.83
E3k.1	0.67	425631	10.54	u1432	0.29	224	2.42
E3k.2	0.74	342370	9.41	fl1577	1.66	6705	38.19
E3k.3	0.67	147135	10.30	d1655	0.94	263	6.51
E3k.4	0.73		8.07	vm1748	1.35	2224	4.43
Random Clustered Euclidean			u1817	0.90	449231	5.01	
C1k.0	0.54	337	9.83	rl1889	1.55	10023	11.45
C1k.1	0.41	534	10.84	d2103	1.44	-	8.19
C1k.2	0.42	320	8.79	u2152	0.62	45205	8.10
C1k.3	0.53	214	7.63	u2319	0.02	7068	3.16
C1k.4	0.58	768	9.36	pr2392	1.22	117	5.75
C1k.5	0.58	139	9.29	pcb3038	0.81	80829	7.26
C1k.6	0.73	1247	7.07	fl3795	1.04	69886	123.66
C1k.7	0.58	449	13.24	fnl4461	0.55	-	12.47
C1k.8	0.34	140	10.40	rl5915	1.56	-	42.00
C1k.9	0.66	703	9.61	rl5934	1.38		56.15
C3k.0	0.62	16009	53.03	pla7397	0.58	-	55.42
C3k.1	0.61	17754	126.49	rl11849	1.02		102.41
C3k.2	0.70	18237	80.39	usa13509	0.66	-	120.20
C3k.3	0.57	6349	71.57	d15112	0.52		90.13
C3k.4	0.57	4845	44.02				
			Randor	n Matrices			
M1k.0	0.01	60	5.47	M3k.0	0.00	612	40.35
M1k.1	0.03	137	5.51	M3k.1	0.01	546	39.52
M1k.2	0.01	151	5.63	M10k.0	0.00	1377	367.84
M1k.3	0.01	169	5.26				

How strong is the Subtour LP bound?

- What about in theory?
- Define

SUBT(c) as the optimal value of the Subtour LP for costs c

- OPT(c) as the length of the optimal tour for costs c
- C_n is the set of all symmetric cost functions on n vertices that obey triangle inequality.
- Then the integrality gap of the Subtour LP is

$$\gamma \equiv \sup_{n} \gamma(n)$$
 where $\gamma(n) \equiv \sup_{c \in \mathcal{C}_n} \frac{OPT(c)}{SUBT(c)}$

A lower bound

It's known that $\gamma \ge 4/3$, where c(i,j) comes from the shortest *i*-*j* path distance in a graph G (graph TSP).



Christofides' Algorithm

Christofides (1976) shows how to compute a tour in polynomial time of cost 3/2 optimal: compute a min-cost spanning tree, compute a matching on the odd-degree vertices, then "shortcut" a traversal of the resulting Eulerian graph.



An upper bound

- Wolsey (1980) and Shmoys and W (1990) show that OPT(c) can be replaced with SUBT(c), so that Christofides gives a tour of cost ≤ 3/2 SUBT(c).
- Therefore,

$$OPT(c) \leq \frac{3}{2}SUBT(c) \Rightarrow \gamma \leq \frac{OPT(c)}{SUBT(c)} \leq \frac{3}{2}$$

Perfect Matching Polytope

Edmonds (1965) shows that the min-cost perfect matching can be found as the solution to the linear program:

Matchings and the Subtour LP

Then MATCH(c) $\leq 1/2$ SUBT(c) since z = 1/2 x is feasible for the matching LP.

 $\begin{array}{ll} \text{Minimize } \sum_{e \in E} c(e)x(e) & \text{Minimize } \sum_{e \in E} c(e)z(e) \\ \text{subject to } & \text{subject to } \sum_{e \in \delta(v)} z(e) = 1 & \forall v \in V \\ & \sum_{e \in \delta(v)} x(e) = 2 & \forall v \in V & \sum_{e \in \delta(S)} z(e) = 1 & \forall S \subset V, |S| \text{ odd} \\ & \sum_{e \in \delta(S)} x(e) \geq 2 & \forall S \subseteq V, |S| \geq 2 \end{array}$

 $0 \le x(e) \le 1 \qquad \forall e \in E$

Shmoys and W (1990) also show that SUBT(c) is nonincreasing as vertices are removed so that matching on odd-degree vertices is at most 1/2 SUBT(c).

Spanning Tree Polytope

Similarly, Edmonds (1971) showed that the mincost spanning tree can be found as the solution of the following LP:

$$\begin{array}{l} \text{Minimize} \sum_{e \in E} c(e) z(e) \\ \text{subject to} \sum_{e \in E} z(e) = |V| - 1 \\ \\ \sum_{e \in E(S)} z(e) \leq |S| - 1 \quad S \subset V \end{array}$$

Spanning Trees and the Subtour LP

Then MST(c) \leq ((n-1)/n) SUBT(c) since z = ((n-1)/n) x is feasible for the MST LP.

$$\begin{array}{ll} \text{Minimize} & \sum_{e \in E} c(e)x(e) \\ & \text{subject to} \\ & \sum_{e \in \delta(v)} x(e) = 2 \quad \forall v \in V \\ & \sum_{e \in E(S)} x(e) \leq |S| - 1 \quad \forall S \subset V, |S| \geq 2 \\ & 0 \leq x(e) \leq 1 \quad \forall e \in E \end{array} \qquad \begin{array}{ll} \sum_{e \in E} \frac{n}{n} \sum_{e \in E} x(e) \\ & \sum_{e \in E(S)} \frac{n}{e(e)} \sum_{e \in E} n \\ & \sum_{e \in E(S)} \frac{n}{e(e)} \sum_{e \in E} n \\ & \sum_{e \in E(S)} \frac{n}{e(e)} \sum_{e \in E} x(e) \\ & = \frac{n-1}{n} \sum_{e \in E(S)} \frac{n}{e(e)} \sum_{e \in E} x(e) \\ & = \frac{n-1}{n} \cdot \frac{1}{2} \cdot 2n \\ & = n-1 \end{array}$$

1

Recent results

- Some recent progress on graph TSP (costs c(i,j) are the shortest i-j path distances in unweighted graph):
 - Boyd, Sitters, van der Ster, Stougie (2010); Aggarwal, Garg, Gupta (2011): Gap is at most 4/3 if graph is cubic.
 - Oveis Gharan, Saberi, Singh (2010): Gap is at most 3/2 ε for a constant ε > 0.
 - Mömke, Svensson (2011): Gap is at most 1.461.
 - Mömke, Svensson (2011): Gap is 4/3 if graph is subcubic (degree at most 3).
 - Mucha (2011): Gap is at most $13/9 \approx 1.44$.
 - Sebő and Vygen (2012): Gap is at most 1.4.

Recent results

- Also some exciting progress on asymmetric TSP (c(i,j) \neq c(j,i)) with triangle inequality (for related LP):
 - Frieze, Galbiati, Maffioli (1982): Gap is O(log n)
 - Asadpour, Goemans, Madry, Oveis Gharan, Saberi, Singh (2010): Gap is O(log n/log log n)
 - Anari, Oveis Gharan (2016): Gap is O(poly log log n) (existential)
 - Svensson, Tarnawski, Vegh (August 2017): Gap is constant
 - ≈ 5500
 - Charikar, Goemans, Karloff (2006): Gap is at least 2

Current state

$$\frac{4}{3} \le \gamma \le \frac{3}{2}$$

• Conjecture (Goemans 1995, others): $\gamma = \frac{4}{3}$

More ignorance

We don't even know the equivalent worstcase ratio between 2-matching costs 2M(c) and SUBT(c).

$$\mu \equiv \sup_{n} \mu(n) \text{ where } \mu(n) \equiv \sup_{c \in \mathcal{C}_{n}} \frac{2M(c)}{SUBT(c)}$$

Then all we know is that

$$\frac{10}{9} \le \mu \le \frac{4}{3}$$
 (Boyd, Carr 1999)

Conjecture (Boyd, Carr 2011): $\mu = \frac{10}{9}$

Our contributions

 We can prove the Boyd-Carr conjecture (with Schalekamp and van Zuylen)

Outline

- A brief intro to the TSP
- A standard TSP linear program
 - Experimental analysis
 - Theoretical analysis: an outstanding open question
- A related question: the Boyd-Carr conjecture and its proof
 - $\mu \leq 4/3$ under a certain condition.
 - $\mu \le 10/9$.
- Some conjectures and more experiments

Some terminology



The strategy

- Start with an optimal fractional 2-matching; this gives a lower bound on the Subtour LP.
- Add a low-cost set of edges to create a graphical 2matching: each vertex has degree 2 or 4; each component has size at least 3; each edge has 0, 1, or 2 copies.



• "Shortcut" the graphical 2-matching to a 2-matching.



Consider fractional 2-matchings that have no cut edge; we show that we can get a graphical 2-matching with a 4/3 increase in cost.



 $2M \leq Graphical 2M \leq 4/3$ Fractional $2M \leq 4/3$ Subtour

Create new graph by replacing path edges with a single edge of cost equal to the path, cycle edges with negations of their cost.



New graph is cubic and 2-edge connected.

Compute a min-cost perfect matching in new graph.



In the fractional 2-matching, double any path edge in matching, remove any cycle edge. Cost is paths + cycles + matching edges.



Why this works

For any given node on the cycle, either its associated path edge is in the matching or one of the two cycle edges.



Why this works

For any given node on the path, either its associated path edge is in the matching or not.



Bounding the cost

- P = total cost of all path edges
- C = total cost all cycle edges
- So fractional 2-matching costs P + C/2
- Claim: Perfect matching in the new graph costs at most 1/3 the cost of all its edges, so at most 1/3(P - C)

Bounding the cost

Since the graphical 2-matching costs at most
 P + C + matching, it costs at most

$$P + C + \frac{1}{3}(P - C) = \frac{4}{3}P + \frac{2}{3}C = \frac{4}{3}\left(P + \frac{1}{2}C\right)$$

 $2M \leq Graphical 2M \leq 4/3$ Fractional 2M

 \leq 4/3 Subtour

Matching cost

- Naddef and Pulleyblank (1981): Any cubic, 2-edgeconnected, weighted graph has a perfect matching of cost at most a third of the sum of the edge weights.
- Proof: Set z(e)=1/3 for all e∈E, then feasible for matching LP.



$$\begin{array}{ll} \text{Minimize } \sum_{e \in E} c(e) z(e) \\ \text{subject to } & \sum_{e \in \delta(v)} z(e) = 1 & \forall v \in V \\ & \sum_{e \in \delta(S)} z(e) \geq 1 & \forall S \subset V, |S| \text{ odd} \end{array}$$

 $3|S| = 2|E(S)| + |\delta(S)|$, so for |S| odd, $|\delta(S)|$ odd.

Proving $\mu \leq 10/9$

- To prove stronger results, we give a polyhedral formulation for graphical 2-matchings.
- For all $i \in V$, create i' and i''
 - i' required: must have degree 2
 - i" optional: may have degree 0 or 2
- For all (i,j)∈E, create edges (i',j'), (i',j"), (i",j')



The formulation

$$\begin{split} &\sum_{e \in \delta(i')} y(e) = 2 \qquad \forall i' \\ &\sum_{e \in \delta(i'')} y(e) \leq 2 \qquad \forall i'' \\ &\sum_{e \in \delta(S) - F} y(e) + |F| - \sum_{e \in F} y(e) \geq 1 \qquad \forall S \subseteq V, F \subseteq \delta(S), F \text{ matching}, |F| \text{ odd} \\ &0 \leq y(e) \leq 1 \qquad \forall e \in E \end{split}$$

Can show that the extreme points of this LP are graphical 2-matchings.

Proving $\mu \leq 10/9$

0



Edmonds (1967)

traveling saleman problem [cf. 4]. I conjecture that there is no good algorithm for the traveling saleman problem. My reasons are the same as for any mathematical conjecture: (1) It is a legitimate mathematical possibility, and (2) I do not know.

A good algorithm is known for finding in any graph

A conjecture

- Conjecture: The worst case for the Subtour LP integrality gap occurs for solutions that are fractional 2-matchings.
- Note: we don't even know tight bounds on γ in this case.

Best-of-Many Christofides'

A conjectured algorithm (Oveis Gharan, Saberi, Singh 2010; An, Kleinberg, Shmoys 2012):

- Solve Subtour LP for x.
- Since ((n-1)/n)x in spanning tree polytope, express ((n-1)/n)x as a convex combination of spanning trees.
- Sample a spanning tree from convex combination, run Christofides' algorithm on it.

Experimental Results

	Std	Max Entropy (Best)	Max Entropy (Ave)	Splitting Off (Best)	Splitting Off (Ave)
TSPLIB	9.56%	3.19%	6.12%	5.23%	6.27%
VLSI	9.73%	5.47%	7.61%	6.60%	7.64%
Graph	12.43%	0.31%	1.23%	0.88%	1.77%

Percentages expressed with respect to cost of an optimal tour

(With Kyle Genova)

Why does this help?

Experimentally, almost all degrees of sampled spanning tree are two. The tree costs more, and matching edges are more expensive, but there are a lot fewer edges in the matching.



Standard Christofides'



Best-of-Many Christofides'



Experimental Results

	Tree		Matching			
	Std	Best of Many	Std	Max Entropy	Splitting Off	
TSPLIB	87.47%	98.57%	31.25%	10.75%	10.65%	
VLSI	89.85%	98.84%	29.98%	12.76%	12.78%	
Graph	79.10%	98.23%	39.31%	4.66%	4.34%	

Percentages expressed with respect to cost of an optimal tour

Analysis

- E[cost of tree] \leq SUBT(c) by construction.
- Would need to show E[cost of matching] \leq (1/2 ϵ) SUBT(c) for some $\epsilon > 0$.

s-t TSP path

Given a fixed start vertex s and end vertex t, find the minimum-cost path from s to t visiting every other vertex exactly once.

- Analog of Christofides': 5/3 (Hoogeveen 1991)
- Lower bound on integrality gap: 3/2
- Best-of-Many Christofides': 1.618 (An, Kleinberg, Shmoys 2012)
- Improved analysis: I.6 (Sebő 2013)
- Improved decomposition of trees:
 - 1.599 (Vygen 2015)
 - I.56 (Gottschalk and Vygen 2015)
 - 3/2 + 1/34 (Sebő and Van Zuylen, April 2016)



"Theory is when we understand everything, but nothing works.

Practice is when everything works, but we don't understand why.

At this station, theory and practice are united, so that nothing works and no one understands why."

Thank you for your attention.